

**Title:** Toxicity in aquatic model species exposed to a temporal series of three different flowback and produced water samples collected from a horizontal hydraulically fractured well.

**Authors:** Erik J. Folkerts<sup>1</sup>, Tamzin A. Blewett<sup>1</sup>, Perrine Delompré<sup>1</sup>, W. Tyler Mehler<sup>1</sup>, Shannon L. Flynn<sup>3</sup>, Chenxing Sun<sup>4</sup>, Yifeng Zhang<sup>4</sup>, Jonathan W. Martin<sup>4</sup>, Daniel S. Alessi<sup>2</sup>, Greg G. Goss<sup>1,5</sup>.

<sup>1</sup>Department of Biological Sciences, University of Alberta, Edmonton, T6G 2E9 Alberta, Canada

<sup>2</sup>Department of Earth and Atmospheric Sciences, University of Alberta, Edmonton, T6G 2E3 Alberta, Canada

<sup>3</sup>School of Natural and Environmental Sciences, Newcastle University, Newcastle upon Tyne, NE1 7RU, UK

<sup>4</sup>Division of Analytical and Environmental Toxicology, University of Alberta, Edmonton, T6G 2E9 Alberta, Canada

<sup>5</sup>National Institute for Nanotechnology, Edmonton, Alberta, T6G 2M9, Canada

## **Abstract**

In the present study, we compared the toxicity and associated chemical characterizations of flowback and produced water (FPW) collected from a single horizontal hydraulically fractured well at different time points during FPW production. Since few studies on whole mixture toxicity related to FPW exist, our aims were to determine both overall toxicity of the FPW mixture in a suite of organisms (*Daphnia magna*, *Lumbriculus variegatus*, *Danio rerio*, and *Oncorhynchus mykiss*) and also determine if toxicity changes depending on variation in FPW chemical properties as a function of time sampled (1.33, 72, and 228 hr FPW samples collected immediately post-well production onset were analyzed in current

study). FPW chemical composition was determined via quadra-pole inductively coupled plasma – mass spectrometry/mass spectrometry (ICP-MS/MS), full-scan high performance liquid chromatography/Orbitrap mass spectrometry (HPLC/Orbitrap-MS), and gas chromatography-mass spectrometry (GC-MS). We observed that FPW sampled later in the production process contained higher ion and total dissolved solids concentrations, whereas the highest concentrations of dissolved organic compounds were observed in the earliest FPW sample analyzed. Toxicity associated with FPW exposure was deemed to be species-specific to a certain extent, but general trends revealed the earliest FPW sampled contained highest toxic potential. Accordingly, we theorize that although the saline conditions of FPW are the foremost toxicological drivers to freshwater organisms, dissolved organics associated with FPW significantly contribute to the overall toxicity of exposed organisms.

**Keywords:** Aquatic Toxicology; Model Species; Hydraulic Fracturing; Organics; Lethality

## ***1. Introduction***

Technological advancements and development in the oil and gas sectors have greatly increased our ability to access world-wide geological reserves of oil and gas previously thought to be unattainable. Implementation of unconventional oil and gas (UOG) recovery processes, in the form of horizontal hydraulic fracturing, is largely responsible for increased production of these resources, and is only expected to become more commonly utilized by industry (Gagnon et al., 2016; Vengosh et al., 2014). UOG operations have significantly expanded in the last decade due to advances in horizontal drilling and completions technology. However, research on environmental impacts of accidental releases is still limited and many gaps in knowledge exist related to the environmental hazards associated to horizontal hydraulic fracturing activities. One such gap pertains to flowback and produced water (FPW), the wastewater by-product that returns to the surface following hydraulic fracturing activities. Although terminology of “flowback” versus “produced water” is subject to operator discretion, FPW collectively is a highly complex mixture composed of original fracturing fluids, compounds from the geological formation itself, saline constituents derived from deep, groundwater reservoirs, and potential daughter

compounds from the transformation of initial fracturing fluid or formation constituents in the high pressure -elevated heat environments of the well-bore-formation interface (Lester et al., 2015; DiGiulio and Jackson, 2016; He et al., 2017a; Hoelzer et al., 2016).

One of the most apparent and immediate hazards posed to the environment from UOG activities and FPW production are surface water contamination events following spills/releases of FPW. In the U.S. alone, 210 billion US gallons ( $7.95 \times 10^8 \text{ m}^3$ ) of FPW was produced from 2005 – 2014 (Kondash and Vengosh, 2015). This is especially concerning as a recent study of the Marcellus and Fayetteville shale formations of the eastern U.S. found that on average, wells drilled for hydraulic fracturing purposes were within ~ 300 m of a freshwater surface body of water, such as streams or rivers (Entekin et al., 2011). Although not as developed as its American counterparts, Canadian UOG development has also been increasing, resulting in larger volumes of FPW produced and increased frequencies of accidental releases to the environment (Alessi et al., 2017; Johnson and Johnson 2012). Thus, understanding effects of FPW releases on the aquatic environment has become essential for government and industry mandates such as spill assessment, clean-up, and remediation protocols. As a result of the paucity of FPW toxicological-related research and high variation in spill mitigation requirements among states and provinces, releases of FPW to surface bodies of water have not historically been considered one of the greatest drivers of FPW environmental risk (Kulander, 2013; Richardson et al., 2013). Furthermore, both Canada and the U.S. do not consider FPW as hazardous waste (Goss et al., 2015). Combining this lack of classification with the decentralized and regionally governed management nature of FPW (Ralston and Kalmbach 2018), policies and mandates dictating FPW spill and remediation strategies are often variable and may not adequately address the hazards posed to the environment. Thus, more conclusive and thorough FPW guideline development among government and industry is a pressing concern. However, to combat the increased risk of FPW environmental impacts, industry operations more commonly implement greater mitigative efforts to assess and reduce risk at the multiple stages of well operations (e.g. pre-development, operational, and well shut-down) by utilization of environmental health and safety regulatory matrices,

more structured and transparent spill reporting and clean-up measures, and more thorough reclamation practices. In Alberta, Canada, this has resulted in a distinct reduction over time in number of reported spills per well (AER Compliance Dashboard) and we assume similar reductions have occurred in other jurisdictions.

Recently, efforts to understand the impact of FPW on aquatic organisms have found that the primary sub-lethal effects observed in organisms following FPW exposure are oxidative and biotransformative stress, negative reproductive impacts, and deleterious effects on the cardio-respiratory system (He et al., 2017b; Blewett et al., 2017a; Blewett et al., 2017b; Folkerts et al., 2017a; Folkerts et al., 2017b). There is also evidence that endocrine disruptive effects may additionally be induced in organisms exposed to FPW (He et al., 2018a; He et al., 2018b). However, references providing strict lethality (measured via varying lethal concentration analyses) toxicological information of FPW on freshwater organisms, as required for guideline development, are still largely unavailable.

One of the complications involved with performing any toxicological study related to FPW is the highly complex and chemically diverse nature of the wastewater (Flynn et al., 2019). Compounding this complexity, the chemical makeup of FPW is highly variable depending on the formation being exploited, initial fracturing fluids used, and well operation characteristics (*e.g.* temperature, pressure, shut-in times, period of flowback) (Kim et al., 2016). Thus, an appreciable amount of work is required to obtain any toxicological knowledge of this wastewater, or of the hazards it poses to the environment. To begin addressing if there are significant toxicological concerns in a legislative and policy-driven context, an evaluation of FPW using standard based whole effluent toxicity protocols (termed WET tests; Chapman, 2000) will provide information about whole-mixture effects on the environment and results subsequently may be included in government and industry protocols and policies.

In the present study, acute FPW toxicity was measured in four different common toxicological model species using samples of FPW collected at different time points during production from a horizontally fractured well in the Duvernay shale play in central Alberta, Canada. These different time

points of sampling cover three operational shifts during the fracturing resource recovery process and FPW produced from these operational phases are hypothesized to contain differing chemical profiles and toxicological potentials, with an increased saline and ion presence predicted in later FPW samples. Early FPW from a producing well is theorized to contain a greater number of initial fracturing additives and may be more toxic to aquatic organisms following exposure. However, FPW samples extending into the production phase of a well (> 120 hrs post well production, as observed in previous studies and those preliminarily made by our group) (Zolfaghari Sharak et al., 2014; Goss et al., 2015) tend to increase drastically in saline content. Among the freshwater aquatic organisms tested, less saline tolerant species (e.g. *Daphnia* and zebrafish) are hypothesized to have greatest sensitivity to FPW, although inter-species responses to FPW exposure are expected to be greatly varied considering the split between invertebrate and vertebrate species and varying lifestyles amongst species tested. Our analyses indicate that temporal variation in FPW chemical characteristics - inorganics, non-targeted organics, and polycyclic aromatic hydrocarbons (PAHs) - can significantly alter toxicity to organisms, and that there is great inter-species variability in lethality responses to FPW. Our chemical characterization and toxicological results further the notion of FPW complexity and serve as a reminder that species-specific differences should be considered for understanding the implications of accidental FPW releases to the environment.

## **2. Materials and Methods**

### **2.1. FPW Samples and Animal Maintenance**

All FPW samples used for analyses were provided by Encana Services Company Ltd. (Calgary, AB, CAN). FPW samples were collected from a single horizontal hydraulically fractured well post-stimulation. Three timepoints for FPW collection (and subsequent analysis) were chosen: a 1 hour, 20 min sample (1.33 hr), a 3-day sample (72 hr), and a 9-day, 12 hour sample (228 hr) (see Table 1). A saline solution matching the major cation and anion concentrations of the most saline FPW sample (228 hr) was also created to control for saline-induced responses (see SI Table 1 for ionic composition information).

A total of four model test species were chosen for toxicological experimentation, including two invertebrate and two vertebrate species. Our two chosen invertebrate species included the freshwater cladoceran, *Daphnia magna* (neonates), and the oligochaete *Lumbriculus variegatus*. Our two vertebrate species were zebrafish (*Danio rerio*; embryos) and rainbow trout (*Oncorhynchus mykiss*; embryos and juveniles). All invertebrate species were cultured and maintained in dechlorinated City of Edmonton (Edmonton, AB, CAN) tap water (moderately hard:  $[\text{Na}^+] = 14.6 \text{ mg/L}$ ,  $[\text{Ca}^{2+}] = 55.9 \text{ mg/L}$ ,  $[\text{Mg}^{2+}] = 15.3 \text{ mg/L}$ ,  $[\text{K}^+] = 2.5 \text{ mg/L}$ , pH: 7.9, conductivity:  $395 \pm 0.5 \text{ }\mu\text{S/cm}$ , dissolved oxygen:  $7.5 \pm 0.5 \text{ mg/L}$ , general hardness: 186 mg/L as  $\text{CaCO}_3$ , salinity: 0 ppt) at  $20 \pm 1 \text{ }^\circ\text{C}$ . Zebrafish embryos (wildtype strain AB) were collected from adult fish housed in 30 L tanks ( $\sim 25$  fish/tank, 14 hr light: 10 hr dark photoperiod) in the University of Alberta zebrafish aquaculture facility filled with zebrafish facility water (pH: 7.4, conductivity:  $553 \text{ }\mu\text{S/cm}$ , temperature:  $28.5 \pm 1 \text{ }^\circ\text{C}$ , dissolved oxygen: 7.3 mg/L, hardness: 174 mg/L as  $\text{CaCO}_3$ , salinity: 0 ppt) treated by a RiOs 100 (reverse osmosis) water purification system. Diploid rainbow trout embryos were collected from the Raven Creek Trout Brood Station (Caroline, AB, CAN), brought to the University of Alberta, and maintained at  $10 \text{ }^\circ\text{C}$  in heath-tray stacks connected to a recirculating water chiller apparatus containing 150 L of dechlorinated City of Edmonton tap water filtered and treated by a bio-ball purification layer (to treat for nitrogenous waste products) and ultra-violet light sterilization. Recirculating water for rainbow trout embryos was constantly aerated and maintained at dissolved oxygen levels of  $8.1 \pm 0.3 \text{ mg/L}$ . Periodic water top-ups to the heath-tray-recirculation system were additionally performed. Juvenile rainbow trout ( $2.38 \pm 0.15 \text{ g}$ ) were maintained in aerated flow-through 400 L tanks filled with dechlorinated City of Edmonton tap water at  $10 \pm 1 \text{ }^\circ\text{C}$ . All animal use was approved by the University of Alberta Animal Care Committee under protocols AUP00001334 and AUP00002352. All previously stated variation was calculated to one standard deviation.

## 2.2. Inorganic FPW Characterization

### 2.2.1. Inductively Coupled Plasma (ICP) -MS/MS Analysis

ICP-MS/MS analysis was used to quantify the concentration of cations, bromide, and total sulfur (Figure 1, SI Table 1). Before analysis all samples were filtered through a 0.2  $\mu\text{m}$  nylon filter membrane and diluted by a factor of 850 for Na analysis and 85 for all other elements with 18 M $\Omega$ ·cm ultrapure water. Diluted 10 ml samples were then acidified with 12  $\mu\text{L}$  using 15.698 N trace metal grade nitric acid. For analysis, an Agilent 8800 Triple Quadrupole ICP-MS (ICP-QQQ) was used with a RF power of 1550 W, a RF reflected power of 18 W, a microMist nebulizer and nickel/copper cones. Samples and external standards were analysed in high matrix mode, in which samples were diluted inline with 8 mL/min argon. Additionally, analysis was performed in MS/MS mode for greater mass resolution and a collision gas reaction cell was used with He gas (3 mL/min), O<sub>2</sub> gas (10% max flow) or H<sub>2</sub> (5 mL/min) gas to overcome matrix interferences (SI Table 2.). Instrumental drift was accounted for using 0.5 ppm solution of indium, which was added to each sample using an inline internal standard addition system. For quality assurance and control, a standard solution was run at the start, middle, and end of each run (every 10 sample for n = 3) to quantify the run precision and percent recovery for each element (SI Table 2).

### ***2.2.2. Anion, total organic carbon and total nitrogen analysis***

Ion chromatography was used to determine the concentration of chloride (Figure 1, SI Table 1). Before analysis, samples were filtered through a 0.2  $\mu\text{m}$  nylon filter membrane and diluted by a factor of 2000 using 18 M $\Omega$ ·cm ultrapure water. Analysis was then performed on a Dionex Ion chromatography DX 600 with a 4mm analytical column (AS9-HC), guard column (AG9-HC), and a 4 mm ASRS Ultra suppressor. For total non-purgeable organic carbon (TOC) and total nitrogen (TN) a subsample of the filter sample was diluted by a factor of five and analyzed using on a TOC analyser (Shimadzu model TOC-V-CHS/CSN). For quality assurance and control for IC a control standard solution was run every five samples (n=4) to quantify the run precision and percent recovery for each element (SI Table 3).

## ***2.3 Organic FPW Characterization***

### ***2.3.1. Extraction of Organics***

For each FPW sample, 150 mL of FPW was vacuum filtered through a glass fiber membrane (90 mm diameter, pore size: 0.4  $\mu$ m), then subsequently freeze-dried for 48 hrs. Approximately 50 mL of the aqueous filtrate was then liquid-liquid extracted using a repeated process of 20 mL of dichloromethane (DCM) and shaking for 3 minutes twice. The combined extracts (40 mL) for each sample were concentrated by nitrogen gas evaporation and reconstituted in 1 mL methanol for High Performance Liquid Chromatography/Orbitrap Mass Spectrophotometry (HPLC/Orbitrap-MS) analysis. The internal standard mix (10 ng of each, Wellington Laboratories, ON, CAN) was spiked into the remaining (unextracted) 100 mL of aqueous filtrate, extracted and evaporated as described above, and reconstituted in 3 mL hexane for further cleanup.

Accelerated solvent extraction (ASE) was used for PAH extraction from particulates in the freeze-dried glass filters sediments. A different, separate glass fiber filter was added at the bottom of the extraction cell, and ~ 2-3 g of florisil (pre-cleaned by DCM) was added, followed by dried sediment filters, to which the internal standard mix was then spiked (10 ng of each). ASE cells were filled with solvent (hexane/DCM 4:1 v:v; Optima™, Fisher Scientific, NH, USA), pressurized to 14 MPa, and heated to 80 °C within 6 min. Pressure and temperature were held for 5 min (static extraction), followed by rinsing with cold solvent (50% of the cell volume) and purging with nitrogen gas for 90 s. This extraction cycle was repeated once more. Approximately 40 mL of total extract was gently concentrated by nitrogen gas evaporation and reconstituted in 3 mL of hexane for further cleanup.

### **2.3.2. High Performance Liquid Chromatography/Orbitrap Mass Spectrophotometry (HPLC/Orbitrap - MS)**

Dissolved organic compounds in FPW aqueous filtrate extracts (section 2.3.1.) were analyzed via HPLC/Orbitrap-MS. Briefly, 10  $\mu$ L of diluted organic extract (equivalent to 50  $\mu$ L of the respective original FPW sample) was used for targeted organic compound identification and semi-quantification using reverse phase LC-Orbitrap-MS. Orbitrap MS was operated in positive electrospray ionization mode, and acquisition was in full scan mode ( $m/z$  100 to 2000) at 2.3 Hz, with resolving power set to 120,000 at



$m/z$  400. Tandem mass analysis of targeted compounds was also performed using collision-induced dissociation or high collision dissociation. Please see He et al. 2018a for further details. The identification of target compounds, as listed in Figure 2, was achieved through accurate mass measurement and tandem mass spectra and later confirmed by the corresponding reference standards. The semi-quantification was achieved by an external linear calibration curve analysis of each of the reference standards. Please see Sun et al. 2019 for a more in-depth account of orbitrap characterization, reference standard chemical concentrations and recoveries, and all other quality assurance and control information.

### **2.3.3. Polycyclic Aromatic Hydrocarbon (PAH) Analysis**

In order to characterize and measure FPW sample PAHs, copper powder and anhydrous sodium sulfate (pre-cleaned with DCM) were added into the 3 mL extracts (section 2.3.1.) and vortexed. Solid phase extraction was used for cleanup of all the sample extracts. A Sep-Pak Silica 6 cc Vac cartridge (1 g; Waters, MA, USA) was conditioned with 5 mL solvent (hexane/DCM 7:3 v:v; Optima™, Fisher Scientific, NH, USA), followed by 5 mL of hexane. Extracts (3 mL) were loaded into cartridges and washed with 4 mL of hexane. PAHs were finally eluted with 5 mL of hexane/DCM 7:3 (v/v) which was subsequently concentrated to 200  $\mu$ L for gas chromatography-mass spectrometry (GC-MS) analysis. Details of PAH standards, internal standard corrected recoveries of the detectable PAHs based on accelerated solvent extraction method (range from 60.0–114%) and liquid-liquid extraction (range from 82.9–179%), and the GC–MS instrumental method have been described in previous work (Zhang et al., 2016). Similarly, details of PAH analyte detection limits can be found in He et al. 2018b.

### **2.4. Lethal Concentration (LC) Analyses and Toxicity Determination**

LC analyses ( $LC_{10}$ ,  $LC_{20}$ , and  $LC_{50}$ ) were performed according to Organisation for Economic Co-operation and Development (OECD) guidelines (OECD-202, 2004; OECD-203, 1992; OECD-236, 2013) with some slight adjustments. Dechlorinated City of Edmonton tap water was used for all exposures and FPW dilutions. Fluorescent lighting on a ratio of 14:10 hr light:dark was applied for all exposures. For 96-hr LC analyses, static-renewal exposures were employed where 50% solution changes were made

every 48 hrs. Dead organisms were immediately removed from exposure containers. A total of seven different FPW concentration dilutions (created by mixing raw FPW with dechlorinated tap water) - including a freshwater control - were used for each LC analysis with six replicates for each individual FPW time-point sample at each dilution being employed to ensure statistical robustness. A series of range-finding tests were performed for all species prior to acute lethal toxicity analyses to ensure proper dilutions of FPW were being used to accurately capture LC values of each FPW sample.

All invertebrate exposures were acute 48-hr, static exposure lethal analyses occurring in 30 mL of FPW containing mediums at  $20 \pm 0.5$  °C in 50 mL glass beakers (10 organisms/beaker). *Daphnia* neonates were immediately collected from adult organisms and exposed within 24 hrs following adult brooding. Adult *Lumbriculus variegatus* were collected from cultures and held overnight before being used in bioassays. Fish (both zebrafish and rainbow trout) embryo lethal analyses occurred over a 96-hr, static-renewal exposure period and used ten embryos/beaker for exposures. Observations every 24 hrs under a stereomicroscope were performed to determine mortality. Zebrafish embryos collected from adult fish were immediately placed into petri dishes containing fresh zebrafish facility water and were subsequently observed under a Leica Zoom 2000 stereomicroscope (Leica Camera Co., GER) to determine embryo fertilization. After confirming fertilization (~ 2 hours post fertilization; hpf) and viable status, embryos were transferred to 100 mL of respective FPW solutions in 250 mL glass beakers maintained at  $25 \pm 0.5$  °C for 96-hr LC<sub>50</sub> analyses.

Rainbow trout embryos (7 – 12 days post fertilization; dpf) removed from the recirculating heath-tray chiller unit were immediately placed into 200 mL of respective FPW solutions in 400 mL beakers maintained at  $10 \pm 1$  °C under constant aeration for LC analyses. Observations every 24 hrs under a stereomicroscope were performed to determine embryo mortality. For juvenile rainbow trout 96-hr LC<sub>50</sub> analyses, static-renewal exposures occurred in 10 L glass tanks containing 8 L of respective FPW solutions under constant aeration and were maintained at  $10 \pm 1$  °C. All tanks were allowed to equilibrate with FPW exposure solutions 24 hrs prior to onset of LC analyses. Fish were fasted for 24 hrs prior to experimentation and were not fed for the duration of the 96-hr LC analysis. For each exposure, 7 fish

were used per tank at each concentration. Observations every 24 hrs were performed to determine fish mortality.

## **2.5. Statistical Analyses and Calculations**

All LC analyses and associated 95% confidence intervals (CI) were calculated using a three-parameter probit model (TRAP; Toxicity Relationship Analysis Program, v1.30a, U.S. EPA). Compared LC values which did not have overlapping 95% CIs were considered significantly different. All graphing and non-linear curve-fitting procedures were performed using the statistic and graphing program Prism (GraphPad Software Inc., CA, USA).

## **3. Results**

### **3.1. Inorganic Characterization of FPW**

ICP-MS/MS inorganic analyses revealed a predicted increasing ion concentration gradient with increased FPW sampling period. This was highlighted in many of the major salt ion characterizations; ions such as Cl, Na, Ca, K, and Mg, which were all shown to increase in concentration as the timing of FPW sampling progressed (Figure 1, SI Table 1). Accordingly, a similar increase in total dissolved solid (TDS) readings was observed, as the highest recorded value was found in the 228 hr sample at ~175700 mg/L. Interestingly, total organic carbon (TOC) values dropped as FPW sampling progressed (SI Table 1), while total nitrogen (TN) measurements did not follow any apparent trend. Regarding metals, high concentrations of Fe, Sr, and Ba were found in all FPW samples, although only Sr displayed a rough observable trend of increasing concentration with FPW sampling time (SI Table 1). Outside these constituents, Br was also found in FPW samples at high concentrations (up to ~ 270 mg/L), but with no obvious trend (SI Table 1).

### **3.2 Organic Characterization of FPW**

Following HPLC/Orbitrap-MS/MS analyses, the organic chemical profiles revealed distinct differences between FPW sampling periods. However, the organic composition of FPW is complex and the lack of reference standards makes quantification difficult, thus, only the organic compounds that were

identified with high confidence (i.e. comparison with corresponding reference standards) are semi-quantified and presented in this study. Polyethylene glycols (PEGs) were found in all FPW samples, with the highest concentrations found in the 1.33 hr sample (63.5 ng/L; Figure 2). Similarly, C<sub>10</sub>-alkyl ethoxylates (C<sub>10</sub>-AEO) and octylphenol ethoxylates (OPE) were present in all FPW samples, with the highest concentration of C<sub>10</sub>-AEO (111 ng/L) found in the 1.33 hr sample and the concentration of OPE highest in the 72 hr FPW sample (85.2 ng/L). Beyond these chemicals, triphenyl phosphate, tri (2-butoxyethyl) phosphate, 3-(dodecanoylamino)-N,N-dimethylpropan-1-amine oxide, and 2-[dimethyl-[3-(tetradecanoylamino)propyl]azaniumyl]acetate were found at varying levels across the FPW samples analyzed, although all were found at very low concentrations overall (Figure 2).

HPLC-MS/MS analyses of 22 PAHs (16 identified by the USA EPA as priority PAHs of toxicological concern; Office of the Federal Registration, 1982) in the aqueous and sediment phases of the FPW samples revealed that all samples contained PAHs; the majority being either of 3 and 4-ringed analytes (Table 2). In particular, the 3-ringed PAH phenanthrene was commonly found at the highest concentrations, with largest total (aqueous and sediment phase summation) concentration of 36.9 µg/L phenanthrene found in the 1.33 hr FPW sample. Phenanthrene (and phenanthrene related compounds such as 1- methylphenanthrene and 3,6-dimethylphenanthrene) were also found to compose the majority of the PAH compounds present in FPW samples, comprising 66%, 63%, and 71% of the 22 total PAHs analyzed in the 1.33 hr, 72 hr, and 228 hr FPW samples, respectively (Table 2). Overall, all PAH compounds followed the general trend of having the highest concentration in the 1.33 hr sample, and lowest in the 72 hr FPW sample, resulting in total PAH concentrations of 112.7, 11.6, and 25.1 µg/L in 1.33 hr, 72 hr, and 228 hr FPW samples, respectively.

### ***3.3. Aquatic Species Lethality Responses***

For clarity and succinctness, all statements of LC analyses and discussion of results will pertain to LC<sub>50</sub> analyses unless otherwise stated. However, information on LC<sub>10</sub> and LC<sub>20</sub> analyses can be found in Table 3. From our inorganic analyses, we anticipated the inherently high salinity of FPW to be a significant contributor to the toxicological hazard of FPW spills posed to freshwater aquatic organisms.

However, our organic chemical characterizations also revealed a substantial amount and diversity of organic compounds to be present. Thus, when compared to our saline control exposures, not all organismal toxic responses to the experimental FPW series samples are observed to be strictly implicated to the high salt concentrations of the FPW samples.

*Daphnia magna* responses to our FPW samples were observed to be most sensitive of the four species tested and displayed the greatest acute toxicity regardless of sample collection timepoint. Comparing the daphnid LC<sub>50</sub> FPW dilution values of 1.58, 0.76, 0.94, and 2.81% for our 1.33 hr, 72 hr, 228 hr, and salt control, respectively (Figure 3, Table 3), to the observed LC<sub>50</sub> values observed in rainbow trout embryos (8.82, 12.26, 10.83, and 16.87% for 1.33 hr, 72 hr, 228 hr, and salt control, respectively) highlights the large variability in aquatic organism response to FPW. Relatively moderate LC<sub>50</sub> responses of zebrafish embryos (3.03, 2.22, 2.15, and 2.11%), *Lumbriculus* (3.13, 4.18, 3.80, and 4.11%), and juvenile rainbow trout (7.14, 10.51, 9.02, and 14.24%) when exposed to the FPW series (1.33 hr, 72 hr, 228 hr, and salt control, respectively) (Figure 3, Table 3) were additionally observed. These results again highlight both the diverse mechanisms and toxic nature of FPW solutions.

#### **4. Discussion**

A significant contribution to the differences in toxicity between the FPW samples analyzed can be attributed to noted differences in inorganic and organic chemical profiles. The FPW samples chosen for analysis from the horizontal hydraulically-fractured well studied spanned a time series hypothesized to hold varying toxicological potentials, as all three time points represented a shift in general well production states.

When individually assessing toxicity amongst species, it was observed that *Daphnia magna* were overall the most sensitive species (lowest LC values) of those tested to FPW samples at all collection time points (Figure 3, Table 3). Interestingly, *Daphnia* were most sensitive to our 72 hr FPW sample when all other species (save for zebrafish embryos) were least sensitive to this particular sample. However, differential *Daphnia* responses to the three FPW samples were insignificant (overlapping LC<sub>50</sub> 95% confidence intervals), but all elicited significantly higher toxic responses (lower LC<sub>50</sub> values with non-

overlapping 95% CIs) compared to the salt controls, indicating that something besides salinity in FPW (presumably organics) causes an additional significant toxic effect. Zebrafish embryos were found to be the second most sensitive to FPW exposures amongst all species tested. However, zebrafish embryo responses generally contrasted other species responses, as the saline control elicited the greatest toxicity in zebrafish embryos and was significantly more toxic compared to the 1.33 hr sample (non-overlapping LC<sub>50</sub> 95% CIs). These results suggest that zebrafish embryos are, above all, most sensitive to high osmotic conditions, and that salinity dominated to a greater degree FPW toxicity in zebrafish embryos, perhaps masking any other effects. Considering their current natural ecological niche and no recent evidence of a diadromous lifestyle within the evolution of their species (Engeszer et al., 2007; Imoto et al., 2012), the high sensitivity of zebrafish to hyperosmotic conditions is a logical response. *Lumbriculus* were found to be the third most overall sensitive species to FPW exposure, although general trends of sensitivity in this species were mirrored in both embryonic and juvenile rainbow trout exposures (Figure 3, Table 3). When analyzing these organism response trends to FPW exposure, it was observed that the 1.33 hr sample elicited greatest toxicity, while the 72 hr sample was least toxic. In contrast to zebrafish, the family Salmonidae, of which rainbow trout are a member, have a relatively recent evolutionary history of anadromy (natural transfer between freshwater and seawater environments) (Docker and Heath, 2003; Crespi and Fulton, 2004; Alexandrou et al., 2013). Thus, a more saline-tolerant response to FPW was expected and found for rainbow trout. Interestingly, responses to different FPW samples within each distinct rainbow trout exposure series were significantly different from one another (no 95% C.I. overlap for LC<sub>50</sub> analyses). However, juvenile rainbow trout were observed to be significantly more sensitive to FPW exposures compared to embryonic forms; a result consistent with earlier reports suggesting that embryonic life-stages are often more resistant to environmental stressors than larval stages (immediately following hatching) and juvenile forms (Embry et al., 2010; Woltering, 1984). It is thought that the enveloping chorion *in ovo* acts as a barrier to toxicants and other environmental stressors, thereby protecting the developing organism within the embryo (Embry et al., 2010; Cotelli et al., 1988; Pelka et al., 2017; Denluck et al., 2018).

As demonstrated by the total ion chromatograms of HPLC/Orbitrap-MS and PAH analyses of FPW extracts, there are clear differences in both diversity and abundance of organic compounds among samples (Figure 2, Table 2). Of the semi-quantified compounds identified, class C<sub>10</sub>-AEO non-ionic surfactants were highest in our 1.33 hr sample. This particular class of surfactants have known bioaccumulation potentials (Müller et al., 1999; Cheng et al., 2005), with nonspecific narcosis as the suspected primary toxicological mode of action (Dorn et al., 1997; Müller and Escher, 1999). Another semi-quantified class of surfactants identified were OPEs. These are non-ionic surfactants which may mimic endogenous hormones and cause endocrine disruption upon exposure (White et al., 1994; Nimrod and Benson, 1996). These surfactants are degraded/or metabolized into octylphenols (OPs), which are often more toxic and more persistent in the environment (Ying et al., 2002; Ahel and Giger, 1993). Although the concentrations of C<sub>10</sub>-AEOs were below water quality guidelines of the Canadian Environmental Protection Act (70 µg/L; Environment and Climate Change Canada Environmental Protection Act, 1999) and OPEs in the 1.33 hr sample were lowest amongst the FPW samples, the toxicity of surfactants in the 1.33 hr sample may still play a larger role in producing greater effects compared to the other two FPW samples, particularly when other characteristics of the mixture are taken into account. Specifically, PEGs were found at highest levels in the 1.33 hr sample. Although not inherently toxic to fish individually, PEG physiochemical properties may alter the toxicity of other compounds found in the FPW mixture. In relation to surfactants, PEGs are shown to increase surfactant critical micelle concentrations (CMC), decrease average micelle aggregation (N<sub>agg</sub>), and increase surfactant polydispersity (Nagarajan and Wang, 2000), while surfactants which incorporate PEGs directly into their chemical structure (such as tocopheryl polyethylene glycol succinate surfactants; TPGS) increase CMCs and decreased N<sub>agg</sub> values compared to traditional non-ionic surfactants without PEGs associated (Sadoqi et al., 2009). These TPGS surfactants have been shown to be an effective pharmaceutical agent for drug deliveries by increasing drug absorption in tissues (Zhang et al., 2012; Ismailos et al., 1994). Increased PEG presence in our 1.33 hr FPW sample may therefore increase the bioavailability of the surfactants

present in the 1.33 hr sample and allow them to interfere and react to a greater degree with biological surfaces of exposed organisms to induce greater toxicity.

PAH analysis of the current samples similarly affirmed TOC analyses depicting the 1.33 sample to contain the greatest amount of organic compounds (Table 2). Of the PAH compounds investigated, all present/detectable forms were highest in concentration in the 1.33 sample, resulting in the highest  $\Sigma 16$  and  $\Sigma 22$  PAHs concentrations. Specifically, phenanthrene-related PAH compounds were highest in the 1.33 hr sample, although in both the 72 and 228 hr samples, these were also the major PAHs identified. PAHs are known to cause numerous toxic effects in exposed organisms. Although PAH toxicity can be species dependent, given that invertebrates are generally considered to exhibit lower PAH metabolic and elimination capabilities, other factors such as developmental stage are also known to influence PAH toxicity (Meador et al., 1995; James, 1989; Varanasi et al., 1989). Regardless, toxicity associated with PAH exposure may come from the parent molecule itself or, as is more often the case, from metabolite intermediates formed during biotransformation processes. These intermediates may include highly reactive epoxides formed during phase I cytochrome P450 (CYP) monooxygenase oxidation steps which may also be transformed into toxic phenol derivatives depending on the original substrate (Buhler and Williams, 1988; Livingstone, 1998). The creation of these intermediate metabolites through CYP systems (and the effects they have on biological systems) are mediated at differing levels depending on the specific PAH by as many as 70 nuclear receptors, including the aryl hydrocarbon receptor (AhR), retinoid X receptor (RXR), and others (Honkakoski and Negishi, 2000; Xu et al., 2005).

Alternatively, PAHs may non-specifically interfere with other cellular processes via nonspecific narcosis (van Wezel and Opperhuizen, 1995; Barron et al., 2004; Di Toro et al., 2007) or other cellular processes/receptors not yet studied. PAHs are also known to produce oxidative stress responses in both invertebrates and vertebrate species (Dalton et al., 2002; Sun et al., 2006; Lemaire et al., 1994; Penning et al., 1996; Zangar et al., 2004). Considering phenanthrene, phenanthrene-related compounds, and other 3-ring PAHs (such as fluorene and dibenzothiophene) in our FPW samples are generally found at the highest concentrations compared to other PAHs, we suspect that significant cardiotoxic potentials are



associated with FPW, as recent research often associates ‘blue-sacs disease’ (a condition characterized by larval craniofacial and tail/spine deformities, as well as pericardial and yolk-sac edemas) (He et al., 2012; Spitsbergen et al., 1991) with exposure to these 3-ring PAHs in larval fish species. Exposure to 3-ring PAHs in fish has recently also been shown to directly alter cardiomyocyte function by specifically affecting  $\text{Ca}^{2+}$  and  $\text{K}^{+}$  currents via blockade of  $\text{K}^{+}$  channel and disruption of sarcolemmal and sarcoplasmic reticulum  $\text{Ca}^{2+}$  cycling (Brette et al., 2017; Brette et al., 2014). Detriments to cardiac function would undoubtedly place higher energetic demands and stress in exposed organisms and may be another toxicological hazard warranting assessment when releases of FPW occur.

## **5. Conclusion**

Research on FPW and wastewaters associated with hydraulic fracturing activities has only recently begun in a manner which assesses the toxicological hazards associated with such fluids. In this study, we aimed to not only determine baseline toxicities of FPW in multiple relevant toxicological model species, but also determine how toxicity (and chemical characterizations) change depending on duration of early stage flowback from a well. These data are of value to the reassessment of risk and remediation strategies/requirements and will overall contribute to a better understanding of the potential impacts of FPW releases. Despite the complexity and variations in FPW chemical and toxicological makeup (dependent on numerous geological and operational factors), understanding basic toxicological and chemical characteristic tendencies will help shape governmental and industrial FPW management policies and hazard assessments.

In the present study, we have determined that FPW toxicity is not only species-dependent but is also influenced by the length of time that a well has been producing FPW. A significant component of the toxicity of FPW to freshwater organisms is mediated by the high salinity and organic make-up (from either initial fracturing fluid constituents and formation derived organics) of the wastewater, with even very high dilutions of FPW inducing toxicity in all species tested. However, all species also demonstrated significantly different toxic responses to at least one of the FPW samples tested separate of saline-related

effects, suggesting that other components of FPW are contributing to overall toxic potentials. In particular, earlier samples of FPW collected closer to the beginning of the flowback period (*e.g.* our 1.33 hr sample) were determined to contain the highest organic contaminant concentrations and were also generally most toxic to the organisms studied. Considering these earlier samples contained overall lower salt concentrations, and metal concentrations that were either lower or equal to later FPW samples, it is assumed that organic contaminants were responsible for most of the added toxicity observed. However, all FPW samples contained significant toxic potential and should be considered hazardous regardless of time post-well stimulation when accidental releases to the environment occur.

### ***Acknowledgements***

This research is funded by the Natural Sciences and Engineering Research Council (NSERC) Collaborative Research and Development Grant CRDPJ 460308 – 14. Additional support was provided by Encana Services Company Ltd. to Drs. Dan Alessi, Greg Goss, and Jon Martin. We would like to thank the University of Alberta Science Animal Support Services (SASS) personnel for their assistance with animal care and maintaining research organisms used throughout this study. We would further like to acknowledge and thank Brian Charles and the staff at the Raven Creek Trout Brood Station for their investment into our research program and supply of rainbow trout embryos. The authors of this present study declare no conflicts of interest.

### ***Literature Cited***

- Ahel, M., Giger, W., 1993. Partitioning of alkylphenols and alkylphenol polyethoxylates between water and organic solvents. *Chemosphere* 26, 1471. doi:10.1016/0045-6535(93)90214-p
- Alberta Energy Regulator (n.d.) Compliance Dashboard (AER). Suite 1000, 250 - 5 Street SW, Calgary, Alberta, T2P 0R4. <http://www1.aer.ca/compliancedashboard/index.html>. Retrieved December 4, 2018.
- Alessi, D.S., Zolfaghari, A., Kletke, S., Gehman, J., 2017. Comparative analysis of hydraulic fracturing wastewater practices in unconventional shale development: Water sourcing, treatment and disposal practices. *Canadian Water Resources Journal* 42, 105–121. doi:10.1002/9781119099987.ch4

457 Alexandrou, M.A., Swartz, B.A., Matzke, N.J., Oakley, T.H., 2013. Genome duplication and multiple  
 458 evolutionary origins of complex migratory behavior in Salmonidae. *Mol Phylogenet Evol* 69, 514–523.  
 459 doi:10.1016/j.ympev.2013.07.026

460 Barron, M.G., Carls, M.G., Heintz, R., Rice, S.D., 2004. Evaluation of fish early life-stage toxicity  
 461 models of chronic embryonic exposures to complex polycyclic aromatic hydrocarbon mixtures.  
 462 *Toxicological Sciences* 78, 60–67. doi:10.1093/toxsci/kfh051

463 Blewett, T.A., Delompré, P.L.M., He, Y., Folkerts, E.J., Flynn, S.L., Alessi, D.S., Goss, G.G., 2017a.  
 464 Sublethal and Reproductive Effects of Acute and Chronic Exposure to Flowback and Produced Water  
 465 from Hydraulic Fracturing on the Water Flea *Daphnia magna*. *Environmental Science & Technology* 51,  
 466 3032–3039. doi:10.1021/acs.est.6b05179

467 Blewett, T.A., Weinrauch, A.M., Delompré, P.L.M., Goss, G.G., 2017b. The effect of hydraulic flowback  
 468 and produced water on gill morphology, oxidative stress and antioxidant response in rainbow trout  
 469 (*Oncorhynchus mykiss*). *Scientific Reports* 7, 46582. doi:10.1038/srep46582

470 Brette, F., Machado, B., Cros, C., Incardona, J.P., Scholz, N.L., Block, B.A., 2014. Crude oil impairs  
 471 cardiac excitation-contraction coupling in fish. *Science* 343, 772–776. doi:10.1126/science.1242747

472 Brette, F., Shiels, H.A., Galli, G.L.J., Cros, C., Incardona, J.P., Scholz, N.L., Block, B.A., 2017. A Novel  
 473 Cardiotoxic Mechanism for a Pervasive Global Pollutant. *Scientific Reports* 7, 41476.  
 474 doi:10.1038/srep41476

475 Buhler, D.R., Williams, D.E., 1988. The role of biotransformation in the toxicity of chemicals. *Aquatic*  
 476 *Toxicology* 11, 19–28. doi:10.1016/0166-445x(88)90004-5

477 Environment and Climate Change Canada, 1999. Canadian Environmental Protection Act: Federal  
 478 environmental quality guidelines. <http://www.ec.gc.ca/ese-ees/default.asp?lang=En&n=164786db-1>.  
 479 Accessed November 21, 2018.

480 Chapman, P.M., 2000. Whole effluent toxicity testing-usefulness, level of protection, and risk assessment.  
 481 *Environmental Toxicology and Chemistry* 19, 3–13. doi:10.1002/etc.5620190102

482 Cheng, H., Kontogeorgis, G.M., Stenby, E.H., 2005. Correlation and Prediction of Environmental  
 483 Properties of Alcohol Ethoxylate Surfactants Using the UNIFAC Method. *Industrial & Engineering*  
 484 *Chemistry Research* 44, 7255–7261. doi:10.1021/ie050096b

485 Cotelli, F., Andronico, F., Brivio, M., Lamia, C.L., 1988. Structure and composition of the fish egg  
 486 chorion (*Carassius auratus*). *Journal of Ultrastructure and Molecular Structure Research* 99, 70–78.  
 487 doi:10.1016/0889-1605(88)90034-1

488 Crespi, B.J., Fulton, M.J., 2004. Molecular systematics of Salmonidae: combined nuclear data yields a  
 489 robust phylogeny. *Mol Phylogenet Evol* 31, 658–679. doi:10.1016/j.ympev.2003.08.012

490 Dalton, T.P., Puga, A., Shertzer, H.G., 2002. Induction of cellular oxidative stress by aryl hydrocarbon  
 491 receptor activation. *Chemico-Biological Interactions* 141, 77–95.

492 Denluck, L., Wu, F., Crandon, L.E., Harper, B.J., Harper, S.L., 2018. Reactive oxygen species generation  
 493 is likely a driver of copper based nanomaterial toxicity. *Environmental Science: Nano* 5, 1473–1481.  
 494 doi:10.1039/c8en00055g

495 Di Toro, D.M., McGrath, J.A., Stubblefield, W.A., 2007. Predicting the toxicity of neat and weathered  
496 crude oil: toxic potential and the toxicity of saturated mixtures. *Environmental Toxicology and Chemistry*  
497 26, 24–36.

498 DiGiulio, D.C., Jackson, R.B., 2016. Impact to Underground Sources of Drinking Water and Domestic  
499 Wells from Production Well Stimulation and Completion Practices in the Pavillion, Wyoming, Field.  
500 *Environmental Science & Technology* 50, 4524–4536. doi:10.1021/acs.est.5b04970

501 Docker, M.F., Heath, D.D., 2003. Genetic comparison between sympatric anadromous steelhead and  
502 freshwater resident rainbow trout in British Columbia, Canada. *Conservation Genetics* 4, 227–231.

503 Dorn, P.B., Rodgers, J.H., Dubey, S.T., Gillespie, W.B., 1997. An assessment of the ecological effects of  
504 a C9–11 linear alcohol ethoxylate surfactant in stream mesocosm experiments. *Ecotoxicology* 6, 275–  
505 292.

506 Embry, M.R., Belanger, S.E., Braunbeck, T.A., Galay-Burgos, M., Halder, M., Hinton, D.E., Léonard,  
507 M.A., Lillicrap, A., Norberg-King, T., Whale, G., 2010. The fish embryo toxicity test as an animal  
508 alternative method in hazard and risk assessment and scientific research. *Aquatic toxicology* (Amsterdam,  
509 Netherlands) 97, 79–87. doi:10.1016/j.aquatox.2009.12.008

510 Engeszer, R.E., Patterson, L.B., Rao, A.A., Parichy, D.M., 2007. Zebrafish in the wild: a review of  
511 natural history and new notes from the field. *Zebrafish* 4, 21–40. doi:10.1089/zeb.2006.9997

512 Entrekin, S., Evans-White, M., Johnson, B., Hagenbuch, E., 2011. Rapid expansion of natural gas  
513 development poses a threat to surface waters. *Frontiers in Ecology* ... 9, 503–511. doi:10.1890/110053

514 Flynn, S. L., von Gunten, K., Warchola, T., Snihur, K., Forbes, T. Z., Goss, G. G., Gingras, M. K.,  
515 Konhauser, K. O., Alessi, D. S., 2019. Characterization and implications of solids associated with  
516 hydraulic fracturing flowback and produced water from the Duvernay Formation, Alberta, Canada.  
517 *Environmental Science: Processes & Impacts*, in press. Submission ID: EM-ART-09-2018-000404.R2

518 Folkerts, E.J., Blewett, T.A., He, Y., Goss, G.G., 2017a. Cardio-respirometry disruption in zebrafish  
519 (*Danio rerio*) embryos exposed to hydraulic fracturing flowback and produced water. *Environ Pollut.*  
520 doi:10.1016/j.envpol.2017.09.011

521 Folkerts, E.J., Blewett, T.A., He, Y., Goss, G.G., 2017b. Alterations to Juvenile Zebrafish (*Danio rerio*)  
522 Swim Performance after Acute Embryonic Exposure to Sub-lethal Exposures of Hydraulic Fracturing  
523 Flowback and Produced Water. *Aquatic Toxicology* 193, 50–59. doi:10.1016/j.aquatox.2017.10.003

524 Gagnon, G.A., Krkosek, W., Anderson, L., McBean, E., Mohseni, M., Bazri, M., Mauro, I., 2016.  
525 Impacts of hydraulic fracturing on water quality: a review of literature, regulatory frameworks and an  
526 analysis of information gaps. *Environmental Reviews* 24, 122–131. doi:10.1139/er-2015-0043

527 Goss, G., Alessi, D., Allen, D., Gehman, J., Brisbois, J., Kletke, S., Sharak, A.Z., Notte, C., Thompson,  
528 D.Y., Hong, K., Prosser, C., 2015. Unconventional Wastewater Management: A Comparative Review and  
529 Analysis of Hydraulic Fracturing Wastewater Management Practices Across Four North American  
530 Basins. *Canadian Water Network* (<http://www.cwn-rceca/>).

531 He, Y., Flynn, S.L., Folkerts, E.J., Zhang, Y., Ruan, D., Alessi, D.S., Martin, J.W., Goss, G.G., 2017a.  
532 Chemical and toxicological characterizations of hydraulic fracturing flowback and produced water. *Water*  
533 Res 114, 78–87. doi:10.1016/j.watres.2017.02.027

534 He, Y., Folkerts, E.J., Zhang, Y., Martin, J.W., Alessi, D.S., Goss, G.G., 2017b. Effects on  
535 Biotransformation, Oxidative Stress, and Endocrine Disruption in Rainbow Trout (*Oncorhynchus mykiss*)  
536 Exposed to Hydraulic Fracturing Flowback and Produced Water. *Environmental Science & Technology*  
537 51, 940–947. doi:10.1021/acs.est.6b04695

538 He, Y., Patterson, S., Wang, N., Hecker, M., Martin, J.W., El-Din, M.G., Giesy, J.P., Wiseman, S.B.,  
539 2012. Toxicity of untreated and ozone-treated oil sands process-affected water (OSPW) to early life  
540 stages of the fathead minnow (*Pimephales promelas*). *Water Res* 46, 6359–6368.  
541 doi:10.1016/j.watres.2012.09.004

542 He, Y., Sun, C., Zhang, Y., Folkerts, E.J., Martin, J.W., Goss, G.G., 2018a. Developmental Toxicity of  
543 the Organic Fraction from Hydraulic Fracturing Flowback and Produced Waters to Early Life Stages of  
544 Zebrafish (*Danio rerio*). *Environmental Science & Technology* 52, 3820–3830.  
545 doi:10.1021/acs.est.7b06557

546 He, Y., Zhang, Y., Martin, J.W., Alessi, D.S., Giesy, J.P., Goss, G.G., 2018b. In vitro assessment of  
547 endocrine disrupting potential of organic fractions extracted from hydraulic fracturing flowback and  
548 produced water (HF-FPW). *Environ Int* 121, 824–831. doi:10.1016/j.envint.2018.10.014

549 Hoelzer, K., Sumner, A.J., Karatum, O., Nelson, R.K., Drollette, B.D., O’Conner, M.P., D’Ambro, E.L.,  
550 Getzinger, G.J., Ferguson, P.L., Reddy, C.M., Elsner, M., Plata, D.L. 2016. Indications of transformations  
551 products from hydraulic fracturing additives in shale-gas wastewater, *Environ. Sci. Technol.* 50, 8036-  
552 8048

553 Honkakoski, P., Negishi, M., 2000. Regulation of cytochrome P450 (CYP) genes by nuclear receptors.  
554 *Biochemical Journal* 347, 321–337. doi:10.1042/0264-6021:3470321

555 Imoto, J.M., Saitoh, K., Sasaki, T., Yonezawa, T., Adachi, J., Kartavtsev, Y.P., Miya, M., Nishida, M.,  
556 Hanzawa, N., 2012. Phylogeny and biogeography of highly diverged freshwater fish species (Leuciscinae,  
557 Cyprinidae, Teleostei) inferred from mitochondrial genome analysis. *Gene* 514, 112–124.  
558 doi:10.1016/j.gene.2012.10.019

559 Ismailos, G., Reppas, C., Macheras, P., 1994. Enhancement of cyclosporin A solubility by d-  
560 alphanatocopheryl-polyethylene-glycol-1000 succinate (TPGS). *European Journal of Pharmaceutical*  
561 *Sciences* 1, 269–271. doi:10.1016/0928-0987(94)90021-3

562 James, M.O., 1989. Biotransformation and disposition of PAH in aquatic invertebrates, in: Varanasi, U.  
563 (Ed.), *Metabolism of Polycyclic Aromatic Hydrocarbons in the Aquatic Environment*. CRC Press, Boca  
564 Raton, FL, pp. 66–91.

565 Johnson, E.G., and Johnson, L.A. 2012. Hydraulic fracture water usage in northeast British  
566 Columbia: Locations, volumes and trends. In *Geoscience Reports 2012*, British Columbia  
567 Ministry of Energy and Mines, Victoria, BC. [AER](#)

568  
569

570 Kim, K., Ozbek, P.O., Dhanasekar, A., Prior, A., Carlson, K. 2016. Temporal analysis of flowback and  
571 produced water composition from shale oil and gas operations: impact of frac fluid characteristics, *J. Pet.*  
572 *Sci Eng* 147, 202–210

573 Kondash, A., Vengosh, A., 2015. Water Footprint of Hydraulic Fracturing. *Environmental Science &*  
574 *Technology Letters* 2, 276–280. doi:10.1021/acs.estlett.5b00211

575 Kulander, C.S., 2013. Shale oil and gas state regulatory issues and trends. Case Western Reserve Law  
576 Review 63, 1101 - 1141.

577 Lemaire, P., Matthews, A., Förlin, L., Livingstone, D.R., 1994. Stimulation of oxyradical production of  
578 hepatic microsomes of flounder (*Platichthys flesus*) and perch (*Perca fluviatilis*) by model and pollutant  
579 xenobiotics. Archives of Environmental Contamination ... 26, 191–200.

580 Lester, Y., Ferrer, I., Thurman, E.M., Sitterley, K.A., Korak, J.A., Aiken, G., Linden, K.G., 2015.  
581 Characterization of hydraulic fracturing flowback water in Colorado: implications for water treatment. Sci  
582 Total Environ 512-513, 637–644. doi:10.1016/j.scitotenv.2015.01.043

583 Livingstone, D.R., 1998. The fate of organic xenobiotics in aquatic ecosystems: quantitative and  
584 qualitative differences in biotransformation by invertebrates and fish. Comp Biochem Physiol, Part A Mol  
585 Integr Physiol 120, 43–49.

586 Meador, J.P., Stein, J.E., Reichert, W.L., Varanasi, U., 1995. Bioaccumulation of Polycyclic Aromatic  
587 Hydrocarbons by Marine Organisms, in: Ware, G.W. (Ed.), Reviews of Environmental Contamination  
588 and Toxicology. Springer, New York, NY, pp. 79–165. doi:10.1007/978-1-4612-2542-3\_4

589 Müller, M.T., Escher, B.I., 1999. Membrane toxicity of linear alcohol ethoxylates. Environmental  
590 Toxicology and Chemistry 18, 2767–2774. doi:10.1002/etc.5620181217

591 Müller, M.T., Zehnder, A.J.B., Escher, B.I., 1999. Liposome-water and octanol-water partitioning of  
592 alcohol ethoxylates. Environmental Toxicology and Chemistry 18, 2191–2198.  
593 doi:10.1002/etc.5620181011

594 Nagarajan, R., Wang, C.-C., 2000. Theory of Surfactant Aggregation in Water/Ethylene Glycol Mixed  
595 Solvents. Langmuir 16, 5242–5251. doi:10.1021/la9910780

596 Nimrod, A.C., Benson, W.H., 1996. Environmental estrogenic effects of alkylphenol ethoxylates. Crit  
597 Rev Toxicol 26, 335–364. doi:10.3109/10408449609012527

598 OECD Guidelines for the Testing of Chemicals. Guideline 202: *Daphnia* sp., Acute Immobilisation Test,  
599 adopted April 2004. <https://www.oecd-ilibrary.org/>. Accessed September 6, 2018.  
600

601 OECD Guidelines for the Testing of Chemicals. Test No. 203: Fish, Acute Toxicity Test, adopted July  
602 1992. <https://www.oecd-ilibrary.org/>. Accessed September 6, 2018.

603 OECD Guidelines for the Testing of Chemicals. Test No. 236: Fish Embryo Acute Toxicity (FET) Test,  
604 adopted July 2013. Retrieved from <https://www.oecd-ilibrary.org/>. Accessed September 6, 2018.

605 Office of the Federal Registration (OFR) Appendix A: Priority pollutants. 1982. Federal Registration 47,  
606 52309.

607 Pelka, K.E., Henn, K., Keck, A., Sapel, B., Braunbeck, T., 2017. Size does matter - Determination of the  
608 critical molecular size for the uptake of chemicals across the chorion of zebrafish (*Danio rerio*) embryos.  
609 Aquatic Toxicology 185, 1–10. doi:10.1016/j.aquatox.2016.12.015

610 Penning, T.M., Ohnishi, S.T., Ohnishi, T., Harvey, R.G., 1996. Generation of reactive oxygen species  
611 during the enzymatic oxidation of polycyclic aromatic hydrocarbon trans-dihydrodiols catalyzed by  
612 dihydrodiol dehydrogenase. Chemical Research in Toxicology 9, 84–92. doi:10.1021/tx950055s

613 Ralston, J.J., and Kalmbach, J.A., 2018. Regulating under conditions of uncertainty and risk: Lessons  
 614 learned from state regulation of hydraulic fracturing. *Environmental Practice* 20, 68 - 79.  
 615 doi:10.1080/14660466.2018.1444875.

616 Richardson, N., Gottlieb, M., Krupnick, A., Wiseman, H., 2013. The state of state shale gas regulation.  
 617 Resources for the Future Report. Washington, D.C.  
 618 [http://www.rff.org/files/sharepoint/WorkImages/Download/RFF-Rpt-StateofStateRegs\\_Report.pdf](http://www.rff.org/files/sharepoint/WorkImages/Download/RFF-Rpt-StateofStateRegs_Report.pdf).  
 619 Accessed November 21, 2018.

620

621 Sadoqi, M., Lau-Cam, C.A., Wu, S.H., 2009. Investigation of the micellar properties of the tocopheryl  
 622 polyethylene glycol succinate surfactants TPGS 400 and TPGS 1000 by steady state fluorometry. *J*  
 623 *Colloid Interface Sci* 333, 585–589. doi:10.1016/j.jcis.2009.01.048

624 Spitsbergen, J.M., Walker, M.K., Olson, J.R., Peterson, R.E., 1991. Pathologic alterations in early life  
 625 stages of lake trout, *Salvelinus namaycush*, exposed to 2,3,7,8-tetrachlorodibenzo- p-dioxin as. *Aquatic*  
 626 *Toxicology* 19, 41–71. doi:10.1016/0166-445x(91)90027-7

627 Sun, C., Zhang, Y., Alessi, D.S., Martin, J.W., 2019. Nontarget Profiling of Organic Compounds in a  
 628 Temporal Series of Hydraulic Fracturing Flowback and Produced Waters. *Environ Int*, *In submission*.

629 Sun, Y., Yu, H., Zhang, J., Yin, Y., Shi, H., Wang, X., 2006. Bioaccumulation, depuration and oxidative  
 630 stress in fish *Carassius auratus* under phenanthrene exposure. *Chemosphere* 63, 1319–1327.  
 631 doi:10.1016/j.chemosphere.2005.09.032

632 Varanasi, U., Stein, J.E., Nishimoto, M., 1989. Biotransformation and Disposition of Polycyclic Aromatic  
 633 Hydrocarbons (PAH) in Fish, in: Varanasi, U. (Ed.), *Metabolism of Polycyclic Aromatic Hydrocarbons in*  
 634 *the Aquatic Environment*. CRC Press, Boca Raton, FL, pp. 93–149.

635 Vengosh, A., Jackson, R.B., Warner, N., Darrah, T.H., Kondash, A., 2014. A Critical Review of the Risks  
 636 to Water Resources from Unconventional Shale Gas Development and Hydraulic Fracturing in the United  
 637 States. *Environmental Science & Technology* 48, 8334–8348. doi:10.1021/es405118y

638 White, R., Jobling, S., Hoare, S.A., Sumpter, J.P., Parker, M.G., 1994. Environmentally persistent  
 639 alkylphenolic compounds are estrogenic. *Endocrinology* 135, 175–182. doi:10.1210/endo.135.1.8013351

640 Woltering, D.M., 1984. The growth response in fish chronic and early life stage toxicity tests: A critical  
 641 review. *Aquatic Toxicology* 5, 1–21. doi:10.1016/0166-445x(84)90028-6

642 Xu, C., Li, C.Y.-T., Kong, A.-N.T., 2005. Induction of phase I, II and III drug metabolism/transport by  
 643 xenobiotics. *Arch Pharm Res* 28, 249–268.

644 Ying, G.-G., Williams, B., Kookana, R., 2002. Environmental fate of alkylphenols and alkylphenol  
 645 ethoxylates--a review. *Environ Int* 28, 215–226.

646 Zangar, R.C., Davydov, D.R., Verma, S., 2004. Mechanisms that regulate production of reactive oxygen  
 647 species by cytochrome P450. *Toxicology and Applied Pharmacology* 199, 316–331.  
 648 doi:10.1016/j.taap.2004.01.018

649 Zhang, Y., Shotyk, W., Zacccone, C., Noernberg, T., Pelletier, R., Bicalho, B., Froese, D.G., Davies, L.,  
 650 Martin, J.W., 2016. Airborne Petcoke Dust is a Major Source of Polycyclic Aromatic Hydrocarbons in  
 651 the Athabasca Oil Sands Region. *Environmental Science & Technology* 50, 1711–1720.  
 652 doi:10.1021/acs.est.5b05092

653 Zhang, Z., Tan, S., Feng, S.-S., 2012. Vitamin E TPGS as a molecular biomaterial for drug delivery.  
654 Biomaterials 33, 4889–4906. doi:10.1016/j.biomaterials.2012.03.046

655 Zolfaghari Sharak, A., Noel, M., Dehghanpour H., Bearinger, D., 2014. Understanding the origin of  
656 flowback salts: A laboratory and field study. Presented at the SPE/CSUR Unconventional Resources  
657 Conference – Canada, Calgary, Alberta: Society of Petroleum Engineers. doi:10.2118/171647--- MS

658 van Wezel, A.P., Opperhuizen, A., 1995. Narcosis due to environmental pollutants in aquatic organisms:  
659 residue-based toxicity, mechanisms, and membrane burdens. Crit Rev Toxicol 25, 255–279.  
660 doi:10.3109/10408449509089890